

# **Dementias Associated with parkinsonism: diagnosis and management**

**Web Ross, MD**

**Dept. of Veterans Affairs**

**Pacific Health Research and Education Institute**

**UH JABSOM**

**August 16, 2023**

**Dementia Series Workshop  
for Catholic Charities Hawaii**

## Sponsored by



CATHOLIC CHARITIES  
HAWAII  
CIRCLE OF CARE FOR DEMENTIA



PACIFIC HEALTH  
RESEARCH & EDUCATION INSTITUTE



These caregiver workshops are made possible by the Elderly  
Affairs Division of the City & County of Honolulu  
through Federal Older Americans Act funding.



# Disclosures

- Dr. Ross had no financial relationships with commercial interests within the past twelve months
- There are no further disclosures.

# Topics for discussion

- **Define dementia associated with parkinsonism, and review frequency**
- **Discuss the diagnosis of different causes of dementia with parkinsonism**
- **Compare the clinical and pathological features of these dementias**
- **Discuss management of these conditions including cognitive and behavioral symptoms**
- **Discuss the effects of these conditions on caregivers**

# Parkinsonian dementias

- **More common disorders:**
  - Parkinson's disease with dementia
  - Dementia with Lewy bodies
- **Atypical parkinsonism and less common:**
  - Progressive supranuclear palsy
  - Multiple Systems atrophy
  - Cortico-basal degeneration
  - Vascular parkinsonism with dementia

## Dementia:

- **acquired impairment in memory and at least one other cognitive domain including**
  - language,
  - visuospatial ability
  - problem solving ability
  - Mood, personality or judgment
- **Severe enough to interfere with social and / or occupational functioning.**

## parkinsonian syndrome:

**Any 2 of :**

**Rest tremor**

**Rigidity or joint stiffness**



**Bradykinesia or slow movement**

**Postural instability or imbalance**

# Frequency of parkinsonian dementias

- Parkinson's disease is present in approximately 1% US population over the age of 65
  - 30 to 40 % of these have dementia
- Dementia with Lewy bodies may account for 25% of all cases in dementia specialty clinic. Prevalence (total number of cases at any given time) is estimated to be 5% over the age of 65 years.
- Atypical parkinsonism prevalence:
  - Multiple system atrophy: between 4 and 5 per 100,000
  - Progressive supranuclear palsy: about 6 per 100,000
  - Corticobasal degeneration: 4 - 5 per 100,000

# Neuropathology in Elderly Decedents with Dementia HAAS vs Rush Aging





# Risk factors:

## PD dementia and dementia with Lewy bodies

- Frequency of both increases with age with average age at onset of 70 to 75 years.
- Both disorders more common in men
- Onset of dementia
  - PD dementia: dementia occurs at least one year and often more than 10 years after onset of the movement disorder
  - Dementia with Lewy bodies: dementia is usually the presenting feature occurring prior to or at the same time as the movement disorder symptoms
- Rapid eye movement sleep behavior disorder may precede both conditions by several years.

# Risk factors:

## REM sleep behavior disorder

- A sleep disorder occurring during the stage of sleep known as rapid eye movement sleep.
- REM sleep is when dreams occur and normally the body is paralyzed during this sleep phase.
- In REM sleep behavior disorder, the normal paralysis is absent and people act out their dreams.
- Dreams may be violent in character and the person may injure their bed partner or themselves.
- REM sleep behavior disorder is present in 40% of PD dementia patients and 90% of DLB patients



# Classic motor features of Parkinson's disease

## Cardinal Features

- Bradykinesia – slow movements
- Rigidity – joint stiffness
- Rest tremor
- Postural instability

## Associated motor Features

- Shuffling gait
- Soft Speech
- Small handwriting
- Masked face



Gowers, 1886

# Parkinson's disease: Non-motor features

- Impaired olfaction
- Constipation
- Heart rate abnormalities
- Excessive daytime sleepiness
- **Rapid Eye Movement sleep behavior disorder**
- Sexual dysfunction
- Fatigue

- Seborrhea
- Dry skin and dry eyes
- Drooling and trouble swallowing
- Bladder dysfunction
- Major depression
- **Dementia**
- **Psychosis**

# PD dementia clinical features

- Movement disorder precedes dementia by at least a year
- Earliest signs of cognitive impairment are in the area of executive function
  - Impaired planning and initiation of complex behaviors such as following directions
  - Trouble switching between tasks or multi-tasking.
  - Impulsive behaviors and lack of ability to monitor self behavior, poor judgement.
- Impaired naming ability, word finding difficulty
- Memory problems: Impaired ability to recall recently learned information

# PD dementia behavioral features

- Hallucinations occur in 45-65% of PD-dementia patients – usually visual
  - Complex and well formed
  - Animals and people
- Depression occurs in about 15% of PD-D
- Apathy or loss of interest is very prominent, occurring in over 50% of PD-D patients.

# Dementia with Lewy bodies: Clinical criteria

- Dementia: acquired and progressive cognitive decline severe enough to interfere with social or occupational functioning.
- Core features (2 required)
  - Fluctuating cognition
  - Recurrent visual hallucinations
  - Parkinsonism

# **Dementia with Lewy bodies: Fluctuating cognition**

- **Marked fluctuating level of consciousness and cognitive abilities that may range from near normal to severe confusion**
  - **Fluctuations may occur over minutes to weeks**
  - **Fluctuations are independent of the normal daily schedule – (not sundowning)**
  - **Person may have episodes of severely reduced levels of arousal (“going blank”) with increased sleepiness**
  - **Attention span very short**



# Dementia with Lewy bodies: Visual Hallucinations

- Recurrent visual hallucinations
  - may be detailed and well formed
  - Frequently of people and animals in the home
  - Response may be indifference, fear, or amusement.
  - Insight may or may not be preserved
  - Worse in cases of visual impairment



# Dementia with Lewy bodies: parkinsonism

- Parkinsonism may occur at the time as dementia or develop later in the course of the condition.
  - bradykinesia and rigidity usually present but milder than Parkinson's Disease.
  - rest tremor usually absent
  - Motor features not necessarily responsive to levodopa



# **Dementia with Lewy bodies: Clinical features of cognitive impairment**

- Short attention span
- Prominent executive function deficits
  - Impaired planning and initiation of complex behaviors like following directions or multi-tasking.
  - Impulsive behaviors and lack of ability to monitor self behavior.
- Prominent visuospatial difficulties or impaired ability to use vision to analyze where objects are in space.
  - Getting lost while walking or driving
- Forgetfulness

# Comparison of early neuropsychological profile


	<b>PD-D</b>	<b>AD</b>	<b>DLB</b>	<b>FTD</b>
<b>Level of consciousness</b>	alert	alert	fluctuates	alert
<b>attention</b>	May be impaired	normal	Severely impaired / fluctuates	normal
<b>Language</b>	Impaired naming	Impaired language early	Variable	Impaired early
<b>memory</b>	forgetfulness	Impaired early	Mild early	Mild early
<b>Visuospatial function</b>	Impaired	impaired	Severely impaired early	Impaired late
<b>Executive function</b>	Severely impaired early	impaired	Severely impaired early	Severely impaired early

# Early clinical profile

	<b>PD-D</b>	<b>AD</b>	<b>DLB</b>	<b>FTD</b>
<b>behavioral symptoms</b>	Visual hallucinations more common	Delusions more common	Visual hallucinations more common	Common, often presenting
<b>rigidity and slow movement</b>	Precede dementia onset	Usually late if at all	May occur with dementia onset	Parkinsonism may occur
<b>rest tremor</b>	May be present	absent	rare	absent

# Pathological features of PD dementia and dementia with Lewy bodies


- Both conditions have Lewy bodies
  - in nerve cells in the brainstem that produce dopamine
  - in the brain cortex
- Both conditions have loss of nerve cells that produce dopamine
- Both may have changes associated with Alzheimer's disease including neuritic plaques and neurofibrillary tangles as well as deficits in the nerve chemical acetylcholine that is involved in memory
- Bottom line: these conditions may not be distinguishable when just looking at the brain.



**Lewy body: target shaped collection of proteins including alpha synuclein within nerve cells that have been injured**



- **Classic**

- Spherical with halo
- Located in brainstem nerve cells that have melanin pigment



- **Cortical**



- Irregular shape, less discrete without halo
- Found in cortical brain regions





**Temporal lobe,  
Synuclein stain, 40X**

# Neuropathology:

- Loss of melanin pigment
- Loss of dopamine producing cells in the substantia nigra
- Lewy bodies
- Low striatal dopamine levels



Lewy bodies in a neuron from the substantia nigra in PD





# Treatment of Dementia with Lewy bodies and PD dementia

- There is no cure and no treatment to slow progression.
- Symptomatic management of dementia includes:
  - Medications
  - Occupational therapy / physical therapy
  - Exercise
  - Management of behavioral complications

# Medications to treat memory problems in people with PD dementia and dementia with Lewy bodies

- Cholinesterase inhibitors – block breakdown of acetylcholine in the brain to help memory
  - Rivastigmine and donepezil associated with modest improvement in cognition
  - May help with anxiety and hallucinations
- Memantine associated with marginal improvement in cognition and behavioral symptoms, but no improvement in activities of daily living

# **Physical, Occupational, and speech / language therapy to assist with management of PD dementia and dementia with Lewy bodies**

- **PT and OT may help with motor symptoms of these conditions**
  - Exercises to maintain range of motion and balance
  - Home modifications to prevent falls and ease daily tasks
- **Use of devices such as canes and walkers may help balance problems and prevent falling**
- **Use of shower chairs and hand rails in the shower and on the commode prevent falls and ease caregiving**
- **Speech therapy:**
  - To assist with maintaining good communication with speech exercises
  - To assist with prevention of aspiration when swallowing

# **management of hallucinations in people with PD dementia and dementia with Lewy bodies**

- **Review medications carefully for meds that can cause hallucinations and decrease dose or discontinue to the extent possible**
  - Dopamine medications that are used to treat the movement disorder such as levodopa, amantadine, pramipexole, ropinirole
  - antidepressants
  - Diphenhydramine
- **Maximize visual acuity**
  - Corrective lenses or cataract surgery
  - Good lighting in hallways and bedrooms including a nightlight
- **Reassurance, comfort, distraction, and safety**

# Medications to manage hallucinations in people with PD dementia and dementia with Lewy bodies

- Work closely with the health care provider, preferably one having expertise in managing hallucinations to
  - Determine if medical therapy is needed
    - Are the hallucinations bothering the patient?
    - Is the patient acting on the hallucinations?
    - Does the patient also have delusions or firmly held ideas that are almost certainly not true?
  - Set goals for therapy
  - Set limits on duration of therapy
- Low doses of atypical antipsychotic medications may be helpful
  - Clozapine
  - Quetiapine
  - Pimavanserin

# Exercise

- Physical exercise improves motor function and balance for people with PD.
- Aerobic exercise (walking, swimming) is better than resistance (weight) training for balance, gait, and overall motor function.
- Tai chi and dance improve balance and tai chi reduces falls up to 6 months after training
- Choose exercise that the person is capable of performing



# Barriers to exercising for persons with PD

- **Low expectation for positive benefit.**
  - **Nearly every form of exercise has been found to be beneficial for PD symptoms and quality of life.**
- **Concern regarding the amount of time needed per day to exercise**
  - **30 minutes of exercise a day is beneficial**
- **The fall risk associated with exercise**
  - **There are many forms of exercise that are safe and that can be performed with a partner**

# Parkinsonian dementias

- **Atypical parkinsonism and less common:**
  - Progressive supranuclear palsy
  - Multiple Systems atrophy
  - Cortico-basal degeneration
  - Vascular parkinsonism with dementia



# Atypical parkinsonism

- **Bradykinesia, rigidity, postural instability are core features**
- **Presence of other clinical features that are not typically seen in PD dementia or dementia with Lewy bodies**
- **Distinct neuropathological abnormalities**
- **Treatment and prognosis of atypical parkinsonism syndromes are different than that of PD**
- **Often misdiagnosed as PD**

# **Atypical parkinsonism dementia features**

- **Mental slowness, apathy, social withdrawal, and fatigue that progresses to dementia with prominent executive function deficits**
- **Behavioral problems including disinhibition, anxiety**
- **Depression**

# Multiple System Atrophy

- Uncommon, neurodegenerative disease associated with parkinsonism without rest tremor
- Onset age is younger than Parkinson's disease (mean = 53 years)
- Balance problems and falls occur early

# **Multiple System Atrophy autonomic features (97%)**

- **Postural dizziness and recurrent fainting**
- **Urinary incontinence early in the disease course**
- **Male impotence may be the presenting feature and is very common**
- **Abnormal sweating (diminished or excessive)**
- **Cold, dusky hands (Raynaud phenomenon)**

# **Multiple System Atrophy non-motor features**

- **REM sleep behavior disorder in 2/3**
- **Sleep disordered breathing including obstructive sleep apnea, central sleep apnea, and stridor**
- **Inspiratory laryngeal stridor**

# Multiple System Atrophy Treatment

- Levodopa and dopamine agonists helpful in about 1/3 at least early in the course for the parkinsonism
- Physical, occupational, and speech therapy

# Multiple System Atrophy Treatment

- **For postural dizziness:**
  - increase salt and water intake
  - elastic leg stockings
  - elevate head of bed at night
  - Frequent small meals
  - caution after meals or exercise
  - Medications
    - Fludrocortisone
    - Midodrine 10 mg tid
    - droxidopa

# **Progressive supranuclear palsy**

- **Underdiagnosed: 1 of 5 are diagnosed in the U.S.**
- **Mean onset age 65 years**
- **Survival is 6 to 10 years after diagnosis**



## **PSP: Early symptoms**

- **Falling may be presenting feature in 2/3**
- **Difficulty walking and imbalance**
- **Slow movement**
- **double and blurred vision**
- **difficulty with speech and swallowing.**

# **PSP: Clinical features**

- **Eye involvement:**
  - **Slow eye movements**
  - **Limited ability to look up or down at first then left and right**
  - **Blepharospasm or forced eyelid closure**
- **Dysphagia or difficulty swallowing**
- **Insomnia**

# **PSP: Treatment**

- **Some respond to dopaminergic medication but good response is short-lived**
- **Amantadine is helpful in some**
- **Botulinum toxin injection for blepharospasm**
- **Physical, occupational therapy**
- **speech therapy for dysarthria and dysphagia**

# Corticobasal degeneration

- **Uncommon**
- **Onset usually after age 60**
- **Men and women equally affected**
- **Survival is 7.9 years after diagnosis**

# **Corticobasal degeneration**

## **Clinical features**

- **parkinsonian syndrome**
- **Cortical sensory loss – inability to correctly interpret sensory information**
- **Action / postural tremor that is jerking**
- **Prominent dysarthria or speech slurring and apraxia of speech (effortful and halting speech)**
- **Slow eye movements**
- **Alien limb**

# **Corticobasal degeneration Treatment**

- **Levodopa usually ineffective**
- **Physical, occupation and speech therapy are helpful**
- **Clonazepam helps the action tremor and myoclonus**

# **Vascular parkinsonism: Clinical features**

- **Vascular risk factors**
  - High blood pressure
  - Diabetes
  - Smoking
  - High cholesterol
- **Lower body parkinsonism:**
  - lower extremity rigidity
  - shuffling gait with start hesitation
  - postural instability
- **Tremor is rare**

# Neuropathological features

- **Alpha synucleinopathies (Lewy bodies)**
  - Parkinson's disease
  - Dementia with Lewy bodies
  - Multiple System Atrophy
- **Tauopathies (neurofibrillary tangles)**
  - Progressive supranuclear palsy
  - Corticobasal degeneration
- **Cerebrovascular lesions (strokes)**
  - Vascular parkinsonism



# Features of parkinsonism dementia most likely to contribute to care provider distress

- Psychosis - Hallucinations, delusions and agitation
- Affective disturbances: Anxiety, depression, and apathy
- Impaired ability to perform activities of daily living such as managing finances, shopping, transportation, and housekeeping; as well as feeding, grooming, dressing, bathing and toileting

**Caregiver burden for parkinsonism dementia is as severe or perhaps more severe than that for Alzheimer's disease**

# Acknowledgements

- **Thank you for your attention!**
- **Thank you to Catholic Charities of Hawaii for inviting me to speak on parkinsonism dementias**
- **Thank you to the Department of Veterans Affairs, Pacific Health Research and Education Institute, Kuakini Medical Center and the University of Hawaii John A Burns School of Medicine for their support.**